

On the Motion of Planets Around the Sun According to Weber's Electrodynamic Law

F. Tisserand

Editor's Note: English translation of Tisserand's 1872 paper: "Sur le mouvement des planètes autour du Soleil, d'après la loi électrodynamique de Weber".¹

Posted in February 2021 at www.ifi.unicamp.br/~assis

¹[Tis72] with English translation in [Tis17a].

1 Editor's Introduction to Tisserand's Paper

By A. K. T. Assis²

Here we present the English translation of Tisserand's 1872 paper: "Sur le mouvement des planètes autour du Soleil, d'après la loi électrodynamique de Weber".³ It has been translated by D. H. Delphenich.⁴ François Félix Tisserand (1845-1896) was a French astronomer.

Wilhelm Weber studied the two-body problem with his force law of 1846.⁵ Weber mentioned Tisserand's paper which is being presented here in his work written in the 1880's and published posthumously in 1894.⁶

Many scientists considered the two-body problem utilizing Weber's law applied to electrodynamics and gravitation.⁷

²Homepage: www.ifi.unicamp.br/~assis

³[Tis72] with English translation in [Tis17a].

⁴feedback@neo-classical-physics.info and <http://www.neo-classical-physics.info/index.html>.

⁵[Web46] with a partial French translation in [Web87] and a complete English translation in [Web07]; see especially [Web71] with English translation in [Web72].

⁶[Web94, Section 3] with English translation in [Web08, Section 3].

⁷[See64] with German translation in [See24], see also [Nor65, p. 46]; [Hol70] with English translation in [Hol17]; [Tis72] with English translation in [Tis17a], [Tis90] with English translation in [Tis17b], see also [Tis96, Volume 4, Chapter 28 (Vitesse de propagation de l'attraction), pp. 499-503] and [Poi53, pp. 201-203]; [Zöl72, p. 334], [Rie74]; [Zöl76b, pp. xi-xii], [Zöl76a, p. 216] and [Zöl83, pp. 126-128]; [Lol83]; [Ser85]; [Rit92]; [Sch97]; [Ger98] with English translation in [Ger], and [Ger17]; [Zen21, pp. 46-47]; [See17a] and [See17b]; [Sch25] with Portuguese translation in [XA94] and English translation in [Sch95]; [Bus26]; [Wie60], [Whi73, pp. 207-208], [Eby77], [SS87], [Ass89], [CA91], [Ass92], [AC92], [Ass94], [Ass99], [AW03], [AWW11], [Ass13], [AWW14], [Ass14], [Ass15], [AWW18] and [FW19].

2 On the Motion of Planets Around the Sun According to Weber's Electrodynamic Law

By F. Tisserand^{8,9,10}

Under that law, the force that produces the motion of the planet around the Sun is:

$$F = \frac{fm\mu}{r^2} \left(1 - \frac{1}{h^2} \frac{dr^2}{dt^2} + \frac{2}{h^2} r \frac{d^2r}{dt^2} \right),$$

in which f is the constant of universal attraction, m is the mass of the planet, μ the sum of that mass and that of the Sun, r is the distance from the planet to the Sun, and h is the velocity by which the attraction propagates in space.

The integration of the equations of motion is accomplished rigorously with the aid of elliptic function. Upon starting with that solution, one can obtain some approximate formulas that will be convenient for the sake of obtaining numerical values. Nonetheless, one will arrive at the goal more rapidly by setting:

$$F = \frac{fm\mu}{r^2} + F_1$$

and regarding F_1 as a perturbing force. Moreover, it will suffice to vary the constants of the elliptical motion.

Here are the equations of the perturbed motion:

$$\left. \begin{aligned} \frac{d^2x}{dt^2} + \frac{f\mu x}{r^2} + X &= 0, \\ \frac{d^2y}{dt^2} + \frac{f\mu y}{r^2} + Y &= 0, \\ \frac{d^2z}{dt^2} + \frac{f\mu z}{r^3} + Z &= 0, \end{aligned} \right\} \quad (1)$$

in which

$$\begin{aligned} X &= \frac{f\mu}{h^2} \frac{x}{r^2} \Omega; \\ Y &= \frac{f\mu}{h^2} \frac{y}{r^2} \Omega; \end{aligned}$$

⁸[Tis72] with English translation in [Tis17a].

⁹Translated by D. H. Delphenich, feedback@neo-classical-physics.info and <http://www.neo-classical-physics.info/index.html>. Edited by A. K. T. Assis, www.ifi.unicamp.br/~assis

¹⁰The Notes by A. K. T. Assis are represented by [Note by AKTA:].

$$\begin{aligned} Z &= \frac{f\mu}{h^2} \frac{z}{r^2} \Omega ; \\ \Omega &= -\frac{dr^2}{dt^2} + 2r \frac{d^2r}{dt^2} . \end{aligned}$$

The equations of elliptic motion are obtained by setting $X = Y = Z = 0$ in equations (1). Suppose that these equations have been integrated and let the elliptic elements be taken to be: a , the semi-major axis, e , the eccentricity, φ , the inclination, θ is the longitude of the node, ϖ is that of the perihelion, and ε is that of the epoch. For the present case, one will have formulas that determine the variation of the constants by taking the well-known formulas and replacing the derivative dR/dp of the perturbing function with respect to an arbitrary element p with $X \frac{dx}{dp} + Y \frac{dy}{dp} + Z \frac{dz}{dp} = R_p$ in them.

Now, one has:

$$R_p = \frac{f\mu}{h^2} \frac{\Omega}{r^3} \left(x \frac{dx}{dp} + y \frac{dy}{dp} + z \frac{dz}{dp} \right) = \frac{f\mu}{h^2} \frac{\Omega}{r^2} \frac{dr}{dp} .$$

Since the expression for the radius vector depends upon only a , e , $\varepsilon - \varpi$, one will have:

$$\frac{dr}{d\varphi} = 0 , \quad \frac{dr}{d\theta} = 0 , \quad \frac{dr}{d\varepsilon} = -\frac{dr}{d\varpi} ,$$

and as a result:

$$R_\varphi = 0 , \quad R_\theta = 0 , \quad R_\varepsilon = -R_\varpi .$$

One will easily find the following formulas:

$$\left. \begin{aligned} \frac{da}{dt} &= -\frac{2}{na} R_\varepsilon , & \frac{d\theta}{dt} &= 0 , \\ \frac{de}{dt} &= -\frac{1-e^2}{na^2e} R_\varepsilon , & \frac{d\varpi}{dt} &= -\frac{\sqrt{1-e^2}}{na^2e} R_e , \\ \frac{d\varphi}{dt} &= 0 , & \frac{d\varepsilon}{dt} &= \frac{2}{na} R_a - \frac{\sqrt{1-e^2}}{na^2e} (1 - \sqrt{1-e^2}) R_e . \end{aligned} \right\} \quad (2)$$

One will remark that these formulas (2) that φ and θ are not altered by the perturbing force, which is obvious *a priori*; however, what is less obvious is that the parameter does not change either. Indeed, one will have:

$$\frac{d[a(1-e^2)]}{dt} = -\frac{2}{na} (1-e^2) R_\varepsilon + 2ae \frac{1-e^2}{na^2e} R_\varepsilon = 0 .$$

In order for us to get some idea of the value of the perturbation, we shall develop those perturbations into series that proceed in sines and cosines of

multiples of the mean anomaly ζ and neglect the powers of e that are greater than one.

We first address Ω , which contains the term $r \frac{d^2 r}{dt^2}$; now, we have:¹¹

$$r \frac{d^2 r}{dt^2} = - \left(\frac{dr}{dt} \right)^2 + x \frac{d^2 x}{dt^2} + y \frac{d^2 y}{dt^2} + z \frac{d^2 z}{dt^2} + \frac{dx^2 + dy^2 + dz^2}{dt^2} ,$$

or even, with an approximation that is entirely satisfactory:

$$r \frac{d^2 r}{dt^2} = - \frac{dr^2}{dt^2} - \frac{f\mu}{r} + f\mu \left(\frac{2}{r} - \frac{1}{a} \right) ;$$

it will then result that:

$$\frac{\Omega}{r^2} = 2f\mu \left(\frac{1}{r^3} - \frac{1}{ar^2} \right) - \frac{3}{r^2} \frac{dr^2}{dt^2} ,$$

which is an expression that is developed as follows:

$$\frac{\Omega}{r^2} = n^2 e \left[2 \cos \zeta + \frac{e}{2} (1 + 11 \cos 2\zeta) \right] + \dots ,$$

One will then have:

$$\begin{aligned} R_a &= \frac{f\mu}{h^2} \frac{\Omega}{r^2} \frac{dr}{da} = \frac{2f\mu}{h^2} n^2 e \cos \zeta + \dots , \\ R_e &= \frac{f\mu}{h^2} \frac{\Omega}{r^2} \frac{dr}{de} = - \frac{f\mu}{h^2} n^2 a e \left[1 + \cos 2\zeta + \frac{3e}{4} (3 \cos \zeta + 5 \cos 3\zeta) \right] + \dots , \\ R_\varepsilon &= \frac{f\mu}{h^2} \frac{\Omega}{r^2} \frac{dr}{d\varepsilon} = \frac{f\mu}{h^2} n^2 a e^2 \sin 2\zeta + \dots , \end{aligned}$$

and upon neglecting e^2 , as always, one will deduce that:

¹¹[Note by AKTA:] Due to a misprint this equation appeared in the original as:

$$r \frac{d^2 r}{dt^2} = - \left(\frac{dr}{dt} \right)^2 + x \frac{d^2 x}{dt^2} + y \frac{d^2 y}{dt^2} + z \frac{d^2 z}{dt^2} + \frac{dx^2 + dy^2 + dz^2}{dt^2} .$$

The expression $(dx^2 + dy^2 + dz^2)/dt^2$ should be understood as:

$$\left(\frac{dx}{dt} \right)^2 + \left(\frac{dy}{dt} \right)^2 + \left(\frac{dz}{dt} \right)^2 .$$

$$\begin{aligned}
\frac{da}{dt} &= 0, & \frac{d\theta}{dt} &= 0, \\
\frac{de}{dt} &= -\frac{f\mu ne}{h^2 a} \sin 2\zeta, & \frac{d\varpi}{dt} &= \frac{f\mu n}{h^2 a} \left[1 + \cos 2\zeta + \frac{3e}{4} (3 \cos \zeta + 5 \cos 3\zeta) \right], \\
\frac{d\varphi}{dt} &= 0, & \frac{d\varepsilon}{dt} &= \frac{4f\mu ne}{h^2 a} \cos \zeta,
\end{aligned}$$

so upon integrating those equations:

$$\begin{aligned}
\delta\alpha &= 0, & \delta\theta &= 0, \\
\delta e &= \frac{f\mu e}{h^2 a} \cos 2\zeta, & \delta\varpi &= \frac{f\mu n}{h^2 a} t + \frac{f\mu}{h^2 a} \left[\frac{1}{2} \sin 2\zeta + \frac{9e}{4} \sin \zeta + \frac{5}{4} e \sin 3\zeta \right], \\
\delta\varphi &= 0, & \delta\varepsilon &= \frac{4f\mu e}{h^2 a} \sin \zeta.
\end{aligned}$$

We will then see that the perturbations of the elements are zero or periodic, with the exception of those of $\delta\varpi$, which contain a secular part. Later, we shall confirm that the periodic parts are entirely negligible under the various hypotheses that one can make on the value of h , in such a way that we will arrive at the following conclusion:

Under Weber's law, the elements will remain the same as under Newton's law. Only the longitude of perihelion will be found to have increased by $\frac{f\mu n}{h^2 a} t$, which is a quantity that will get larger as the planet gets closer to the Sun.

Consider the case of Mercury. Upon taking the mean solar day to be the unit of time, and the semi-major axis of the orbit of the Earth to be the unit of distance, one will find that:

$$\delta\varpi = \frac{(1.05160)}{h^2} t.$$

If we assume that h has the same value as in Weber's experiments on electricity, namely, $h = 439450 \times 10^6$, with seconds and millimeters for units, then we will first have:

$$\log h = 2.40805 \qquad \text{and} \qquad \delta\varpi = (\bar{4}.23550)t$$

per century, with our units, and then find that:

$$\delta\varpi = +6.28'' ;$$

for Venus, one will have only:

$$\delta\varpi = +1.32'' .$$

If one supposes that h is equal to the speed of propagation of light, then one will have:

$$\log h = 2.23948$$

and then

For Mercury and one century: $\delta\varpi = +13.65''$,

For Venus and one century: $\delta\varpi = +2.86''$.

In order to show that these periodic terms are negligible, it will suffice to take the biggest of them, which is $\delta\varpi$, namely, $\frac{f\mu}{2ah^2} \sin 2\zeta$; one will find that its coefficient does not reach $0.003''$.

References

- [AC92] A. K. T. Assis and R. A. Clemente. The ultimate speed implied by theories of Weber's type. *International Journal of Theoretical Physics*, 31:1063–1073, 1992.
- [Ass89] A. K. T. Assis. On Mach's principle. *Foundations of Physics Letters*, 2:301–318, 1989.
- [Ass92] A. K. T. Assis. *Curso de Eletrodinâmica de Weber*. Setor de Publicações do Instituto de Física da Universidade Estadual de Campinas — UNICAMP, Campinas, 1992. Notas de Física IFGW Número 5. Available at www.ifi.unicamp.br/~assis and www.bibliotecadigital.unicamp.br/document/?down=60362.
- [Ass94] A. K. T. Assis. *Weber's Electrodynamics*. Springer, Dordrecht, 1994. Doi: 10.1007/978-94-017-3670-1.
- [Ass99] A. K. T. Assis. *Relational Mechanics*. Apeiron, Montreal, 1999. Available at www.ifi.unicamp.br/~assis.
- [Ass13] A. K. T. Assis. *Mecânica Relacional e Implementação do Princípio de Mach com a Força de Weber Gravitacional*. Apeiron, Montreal, 2013. Available at www.ifi.unicamp.br/~assis.
- [Ass14] A. K. T. Assis. *Relational Mechanics and Implementation of Mach's Principle with Weber's Gravitational Force*. Apeiron, Montreal, 2014. Available at www.ifi.unicamp.br/~assis.
- [Ass15] A. K. T. Assis. *Eletrodinâmica de Weber: Teoria, Aplicações e Exercícios*. Editora da Unicamp, Campinas, 2nd edition, 2015.
- [AW03] A. K. T. Assis and K. H. Wiederkehr. Weber quoting Maxwell. *Mitteilungen der Gauss-Gesellschaft*, 40:53–74, 2003.
- [AWW11] A. K. T. Assis, K. H. Wiederkehr, and G. Wolfschmidt. *Weber's Planetary Model of the Atom*, volume 19 of *Nuncius Hamburgensis — Beiträge zur Geschichte der Naturwissenschaften*. Tredition Science, Hamburg, 2011. Edited by G. Wolfschmidt.
- [AWW14] A. K. T. Assis, K. H. Wiederkehr, and G. Wolfschmidt. *O Modelo Planetário de Weber para o Átomo*. Apeiron, Montreal, 2014. Available at www.ifi.unicamp.br/~assis.

- [AWW18] A. K. T. Assis, K. H. Wiederkehr, and G. Wolfschmidt. *Webers Planeten-Modell des Atoms*. Apeiron, Montreal, 2018. Translated by H. Härtel. Available at www.ifi.unicamp.br/~assis.
- [Bus26] V. Bush. The force between moving charges. *Journal of Mathematical Physics*, 5:129–157, 1926.
- [CA91] R. A. Clemente and A. K. T. Assis. Two-body problem for Weber-like interactions. *International Journal of Theoretical Physics*, 30:537–545, 1991.
- [Eby77] P. B. Eby. On the perihelion precession as a Machian effect. *Lettere al Nuovo Cimento*, 18:93–96, 1977.
- [FW19] U. Frauenfelder and J. Weber. The fine structure of Weber’s hydrogen atom: Bohr-Sommerfeld approach. *Zeitschrift für angewandte Mathematik und Physik*, 70:105, 2019. Doi: 10.1007/s00033-019-1149-4.
- [Ger] P. Gerber. The spatial and temporal propagation of gravity. *The General Science Journal*. Translation by H. Jones. Available at <https://www.gsjournal.net/Science-Journals/Historical%20Papers/View/4011> and <http://www.alternativephysics.org/gerber/Perihelion.htm>.
- [Ger98] P. Gerber. Die räumliche und zeitliche Ausbreitung der Gravitation. *Zeitschrift für Mathematik und Physik II*, 43:93–104, 1898.
- [Ger17] P. Gerber. Die Fortpflanzungsgeschwindigkeit der Gravitation. *Annalen der Physik*, 52:415–444, 1917.
- [Hol70] G. Holzmüller. Ueber die anwendung der Jacobi-Hamilton’schen Methode auf den Fall der Anziehung nach dem electrodynamischen Gesetze von Weber. *Zeitschrift für Mathematik und Physik*, 15:69–91, 1870.
- [Hol17] G. Holzmüller, 2017. On the application of the Jacobi-Hamilton method to the case of attraction according to Weber’s electro-dynamical law. Translated by D. H. Delphenich. Available at <http://www.neo-classical-physics.info/index.html>.
- [Lol83] G. Lolling. Ueber Bewegungen elektrischer Theilchen nach dem Weber’schen Grundgesetz der Elektrodynamik. *Verhandlungen der Kaiserlichen Leopoldinisch-Carolinischen Deutschen Akademie der Naturforscher*, 44:273–336, 1883.

- [Nor65] J. D. North. *The Measure of the Universe — A History of Modern Cosmology*. Clarendon Press, Oxford, 1965.
- [Poi53] H. Poincaré. Les limites de la loi de Newton. *Bulletin Astronomique*, 17:121–269, 1953.
- [Rie74] E. Riecke. Ueber Molecularbewegung zweier Theilchen, deren Wechselwirkung durch das Webersche Gesetz der electrischen Kraft bestimmt wird. *Nachrichten von der K. Gesellschaft der Wissenschaften und der Georg-Augustus-Universität*, 27:665–672, 1874.
- [Rit92] E. Ritter. Bewegung eines materiellen mit Elektrizität geladenen Theilchens unter der Einwirkung eines ruhenden Centrums bei Giltigkeit des Weber’schen Gesetzes. *Zeitschrift für Mathematik und Physik*, 37:8–24, 1892.
- [Sch97] W. Scheibner. Ueber die formale Bedeutung des Hamiltonschen Principis und das Weber’sche Gesetz. *Berichte über die Verhandlung der königlich Sächsischen Gesellschaft der Wissenschaften zu Leipzig*, Math.-Phys. Classe, 49:578–601, 1897.
- [Sch25] E. Schrödinger. Die Erfüllbarkeit der Relativitätsforderung in der klassischen Mechanik. *Annalen der Physik*, 77:325–336, 1925.
- [Sch95] E. Schrödinger. The possibility of fulfillment of the relativity requirement in classical mechanics. In J. B. Barbour and H. Pfister, editors, *Mach’s Principle — From Newton’s Bucket to Quantum Gravity*, pages 147–158, Boston, 1995. Birkhäuser. Translated by J. B. Barbour.
- [See64] C. Seegers, 1864. De motu perturbationibusque planetarum secundum legem electrodynamicam Weberianam solem ambientium. Dissertation, Göttingen.
- [See17a] H. Seeliger. Bemerkung zu P. Gerbers Aufsatz: “Die Fortpflanzungsgeschwindigkeit der Gravitation”. *Annalen der Physik*, 53:31–32, 1917.
- [See17b] H. Seeliger. Weitere Bemerkungen zur “Fortpflanzungsgeschwindigkeit der Gravitation”. *Annalen der Physik*, 54:38–40, 1917.
- [See24] C. Seegers. *Über die Bewegung und die Störungen der Planeten, wenn dieselben sich nach dem Weberschen elektrodynamischen*

Gesetz um die Sonne bewegen. Kommissionsverlag von Friedr. Viewweg & Sohn Akt. Ges., Braunschweig, 1924. Neu herausgegeben von P. Heylandt. Übersetzt von F. Diestel.

- [Ser85] A. Servus, 1885. Untersuchungen über die Bahn und die Störungen der Himmelskörper mit Zugrundelegung des Weber'schen electrodynamischen Gesetzes. Dissertation, Halle.
- [SS87] A. G. Sokol'skii and A. A. Sadovnikov. Lagrangian solutions for Weber's law of attraction. *Soviet Astronomy (Astronomical Journal)*, 31:90–93, 1987.
- [Tis72] F. Tisserand. Sur le mouvement des planètes autour du soleil, d'après la loi électrodynamique de Weber. *Comptes Rendues de l'Academie des Sciences de Paris*, 75:760–763, 1872.
- [Tis90] F. Tisserand. Sur les mouvements des planètes, en supposant l'attraction représentée par l'une des lois électrodynamiques de Gauss ou de Weber. *Comptes Rendues de l'Academie des Sciences de Paris*, 110:313–315, 1890.
- [Tis96] F. F. Tisserand. *Traité de Mécanique Céleste*, volume 4, Chapter 28 (Vitesse de propagation de l'attraction), pp. 499-503, "Loi d'attraction conforme à la loi électrodynamique de Weber". Gauthier-Villars, Paris, 1896.
- [Tis17a] F. Tisserand, 2017. On the motion of planets around the Sun according to Weber's electrodynamical law. Translated by D. H. Delphenich. Available at <http://www.neo-classical-physics.info/index.html>.
- [Tis17b] F. Tisserand, 2017. On the motion of planets when one supposes that the attraction is represented by either the electro-dynamical law of Gauss or that of Weber. Translated by D. H. Delphenich. Available at <http://www.neo-classical-physics.info/index.html>.
- [Web46] W. Weber. Elektrodynamische Maassbestimmungen — Über ein allgemeines Grundgesetz der elektrischen Wirkung. *Abhandlungen bei Begründung der Königlich Sächsischen Gesellschaft der Wissenschaften am Tage der zweihundertjährigen Geburtstagfeier Leibnizens's herausgegeben von der Fürstlich Jablonowskischen Gesellschaft (Leipzig)*, pages 211–378, 1846. Reprinted in Wilhelm

Weber's *Werke*, Vol. 3, H. Weber (ed.), (Springer, Berlin, 1893), pp. 25-214.

- [Web71] W. Weber. Elektrodynamische Maassbestimmungen insbesondere über das Princip der Erhaltung der Energie. *Abhandlungen der mathematisch-physischen Classe der Königlich Sächsischen Gesellschaft der Wissenschaften zu Leipzig*, 10:1–61, 1871. Reprinted in Wilhelm Weber's *Werke*, Vol. 4, H. Weber (ed.), (Springer, Berlin, 1894), pp. 247-299.
- [Web72] W. Weber. Electrodynamical measurements — Sixth memoir, relating specially to the principle of the conservation of energy. *Philosophical Magazine*, 43:1–20 and 119–149, 1872. Translated by Professor G. C. Foster, F.R.S., from the *Abhandlungen der mathem.-phys. Classe der Königlich Sächsischen Gesellschaft der Wissenschaften*, vol. x (January 1871).
- [Web87] W. Weber. Mesures électrodynamiques. In J. Joubert, editor, *Collection de Mémoires relatifs a la Physique*, Vol. III: *Mémoires sur l'Électrodynamique*, pages 289–402. Gauthier-Villars, Paris, 1887.
- [Web94] W. Weber. Elektrodynamische Maassbestimmungen insbesondere über den Zusammenhang des elektrischen Grundgesetzes mit dem Gravitationsgesetze. In H. Weber, editor, Wilhelm Weber's *Werke*, Vol. 4, pages 479–525, Berlin, 1894. Springer.
- [Web07] W. Weber, 2007. Determinations of electrodynamic measure: concerning a universal law of electrical action, 21st Century Science & Technology, posted March 2007, translated by S. P. Johnson, edited by L. Hecht and A. K. T. Assis. Available at <http://21sci-tech.com/translation.html> and www.ifi.unicamp.br/~assis.
- [Web08] W. Weber, 2008. Determinations of electrodynamic measure: particularly in respect to the connection of the fundamental laws of electricity with the law of gravitation, 21st Century Science & Technology, posted November 2008, translated by G. Gregory, edited by L. Hecht and A. K. T. Assis. Available at <http://21sci-tech.com/translation.html> and www.ifi.unicamp.br/~assis.

- [Whi73] E. T. Whittaker. *A History of the Theories of Aether and Electricity*, volume 1: *The Classical Theories*. Humanities Press, New York, 1973.
- [Wie60] K. H. Wiederkehr. Wilhelm Webers Stellung in der Entwicklung der Elektrizitätslehre. Dissertation, Hamburg, 1960.
- [XA94] A. L. Xavier Jr. and A. K. T. Assis. O cumprimento do postulado de relatividade na mecânica clássica — uma tradução comentada de um texto de Erwin Schrödinger sobre o princípio de Mach. *Revista da Sociedade Brasileira de História da Ciência*, 12:3–18, 1994. Errata available at www.ifi.unicamp.br/~assis.
- [Zen21] J. Zenneck. Gravitation. In A. Sommerfeld, editor, *Encyklopädie der Mathematischen Wissenschaften mit Einschluss Ihrer Anwendungen*, volume 5, Part 1, chapter 2, pages 25–67. B. G. Teubner, Leipzig, 1903–1921. Abgeschlossen im August 1901.
- [Zöl72] F. Zöllner. *Über die Natur der Cometen. Beiträge zur Geschichte und Theorie der Erkenntnis*. Engelmann, Leipzig, 1872.
- [Zöl76a] F. Zöllner. Ueber die physikalischen Beziehungen zwischen hydrodynamischen und elektrodynamischen Erscheinungen. *Berichte über die Verhandlungen der Königlich Sächsischen Gesellschaft der Wissenschaften zu Leipzig — Mathematisch-physische Classe*, I. II:59–226, 1876.
- [Zöl76b] J. C. F. Zöllner. *Principien einer elektrodynamischen Theorie der Materie*. Engelmann, Leipzig, 1876.
- [Zöl83] J. C. F. Zöllner. *Über die Natur der Cometen — Beiträge zur Geschichte und Theorie der Erkenntnis*. L. Staackmann, Leipzig, 3rd edition, 1883. First edition of 1872.